

Asperger Syndrome



1919 University Avenue West, Suite 400, St. Paul, MN 55104 Tel. 651-645-2948 or 888-NAMIHELPS www.namihelps.org

The *Diagnostic and Statistical Manual of Mental Disorders (4th Edition)*, known as *DSM IV*, published in 1994, defines Asperger syndrome (AS) as marked by "severe and sustained impairment in social interaction" along with "restricted repetitive and stereotyped patterns of behavior."

As a parent, what behaviors might I suspect as indicative of this disorder?

The more general traits that may be observed include awkwardness in social situations, an intense preoccupation with certain specific (often unusual) topics, self-directed orientation, a lack of understanding of social cues, and clumsiness caused by lack of motor coordination.

What is meant by "severe and sustained impairment in social interaction"?

A child with AS often has problems with normally developed verbal as well as non-verbal interaction tools. The child may, for example, not meet the eyes of a person speaking, seem to lack facial expressiveness, or not use normal body posturing and gestures. This affects social interaction in a negative way.

What are "restricted repetitive patterns of behavior"?

This kind of behavior is demonstrated by a preoccupation with certain actions or objects within a restricted range. Rather than applying an intense interest to a variety of subjects, the child with AS has interests of a rather narrow scope, like aliens or computers, bus routes or sports schedules, maps and charts.

This restricted repetitive behavior also is exhibited through a very rigid, non-negotiable adherence to specific nonfunctional routines or rituals. The child with this disorder may, for example, insist on walking a certain route to school without deviation. The child is inflexible about following a certain sequence of events--he or she may need to walk in a circle before sitting down or dress in a specific order. These nonfunctional routines can be of critical importance to the child with Asperger. Given a choice in clothing, the child might create what seems like a uniform that is worn day after day.

How is Asperger different from autism?

A child with Asperger experiences no clinically significant delay in cognitive development and does not experience a gross delay in developing language skills.

Other differences are:

- Children with autism tend to think concretely and have much difficulty with symbolic thinking and pretend play, whereas Asperger children can be quite imaginative although themes may be repetitive.
- Asperger children tend to have motor coordination difficulties not seen in autism.

What about relationships with peers?

The child may not make friends easily, or at all, and may not seem interested in sharing experiences or interests with those around him. For example, a child developing normally may show his artwork to people around him or bring a toy to his sister or brother to look at, but a child with AS will not as readily do so.

A child or adolescent with AS may seem unwilling or uninterested in responding to others in a socially or emotionally reciprocal way. For example, the child may ignore or seem to not notice when a person expresses affection toward him or prompts conversation. On the other hand, Asperger individuals may highly desire social interaction, but their poor social skills result in failure which can cause anxiety and depression.

What about the course of Asperger syndrome?

AS usually presents between ages 2 and 6 years, but is often not recognized until later. As far as doctors know, the disorder is present throughout the course of a person's life. It has often been diagnosed as late as young adulthood.

Who gets Asperger?

Not much is known about how common the syndrome is because few studies have been done. Prevalence rates range from .024 percent to .36 percent based on studies in Canada and Goteborg, Sweden, respectively.

Boys appear to have a higher incidence than girls at a 4:1 ratio. There is likely a genetic component which is thought by some to be related to the genetic deficit in autism. This is presently unclear.

How does one arrive at a diagnosis of Asperger?

The diagnosis is based on the presence of signs and symptoms in the DSM-IV. Differential diagnosis includes autism, complex learning disabilities, schizophrenia-spectrum disorder and obsessive-compulsive disorder.

What treatments can be considered useful or helpful for the child or adolescent with Asperger?

Because securing educational and related services may be difficult due to the lack of knowledge about Asperger, it is important for the parents and clinician to work closely together to supply the patient and school personnel with the necessary information and help. Educational interventions are often necessary and should be individual accommodations to the person's needs. Because these students generally do well with memory tasks, teaching in a rote fashion may help the individual to retain the information presented.

Deficits in social skills may be remediated in small groups usually led by a mental health professional or speech and language pathologist.

Depending on the presence and extremity of associated symptoms, psycho-pharmacological interventions may help. Examples of associated symptoms that may be effectively treated with medication are hyperactivity, impulsivity, inattention, mood instability, temper outbursts, depression, anxiety and obsessive-compulsive symptoms.

Summary:

Early intervention and treatment is the single most important effort a parent can make to influence the outcomes for a child or adolescent with Asperger. Finding a clinician that can make the diagnosis of Asperger may be the more significant hurdle in getting appropriate treatment for your child.

Resources:

Autism and AS, by Uta Frith, Cambridge University Press, London, UK, 1991

For more information on the Internet, look for the "Asperger Disorder Homepage" at

<http://www.ummed.edu/pub/o/ozbayrak/asperger.html>

Updated January 2013; May 2017